

NUPLAZID PRICING AND DISTRIBUTION FACT SHEET

ONCE-DAILY
NUPLAZID[®]
(pimavanserin) 34mg capsules

Overview ¹	
Brand name	NUPLAZID
Generic name	Pimavanserin
National drug codes (NDCs)	NUPLAZID 34 mg capsules (bottle of 30): NDC 63090-340-30 NUPLAZID 10 mg tablets (bottle of 30): NDC 63090-100-30
Indication and usage	NUPLAZID is indicated for the treatment of hallucinations and delusions associated with Parkinson's disease psychosis.
Dosage and administration	The recommended dose of NUPLAZID is 34 mg taken orally once daily, without titration. NUPLAZID can be taken with or without food. NUPLAZID capsules can be taken whole, or opened and the entire contents sprinkled over a tablespoon (15 mL) of applesauce, yogurt, pudding, or a liquid nutritional supplement. Consume the drug/food mixture immediately without chewing; do not store for future use. Coadministration with strong CYP3A4 inhibitors (e.g. ketoconazole) increases NUPLAZID exposure. Reduce NUPLAZID dose to 10 mg taken orally as one tablet once daily. Coadministration with strong or moderate CYP3A4 inducers reduces NUPLAZID exposure. Avoid concomitant use of strong or moderate CYP3A4 inducers with NUPLAZID.
Manufacturer	Acadia Pharmaceuticals Inc.
Distribution	
Specialty distributors	Eligible government, long-term care, and hospital inpatient pharmacies may order NUPLAZID directly from: <ul style="list-style-type: none"> ▪ ASD Healthcare: AmerisourceBergen 1-800-746-6273 ▪ Cardinal Health 1-844-794-7276 ▪ McKesson Plasma and Biologics 1-877-625-2566 ▪ Morris & Dickson 1-800-710-6100
Group purchasing organizations	<ul style="list-style-type: none"> ▪ GeriMed 1-800-456-4374 ▪ MHA 1-800-642-3020
Specialty pharmacy network	<ul style="list-style-type: none"> ▪ Accredo Specialty Pharmacy 1-877-222-7336 ▪ AllianceRx Walgreens Prime 1-888-782-8443 ▪ Humana Specialty Pharmacy 1-800-486-2668 ▪ Advanced Care Scripts 1-877-985-6337 ▪ CVS Specialty Pharmacy 1-800-237-2767 ▪ Walmart Specialty Pharmacy 1-877-453-4566
Price and support	
Wholesale acquisition cost	\$4,173.00 for 30-count bottle (1-month supply) effective February 5, 2022
Support program	Acadia Connect™ provides patient access support, including navigating insurance, managing cost, and filling their prescription. <ul style="list-style-type: none"> ▪ Acadia Connect support hotline: 1-844-737-2223 ▪ Acadia Connect LTC support hotline: 1-877-889-0739

Important Safety Information

WARNING: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

- Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death.
- NUPLAZID is not approved for the treatment of patients with dementia-related psychosis unrelated to the hallucinations and delusions associated with Parkinson's disease psychosis.

Contraindication: NUPLAZID is contraindicated in patients with a history of a hypersensitivity reaction to pimavanserin or any of its components. Rash, urticaria, and reactions consistent with angioedema (e.g., tongue swelling, circumoral edema, throat tightness, and dyspnea) have been reported.

See additional Important Safety Information on the following page. Please read the full [Prescribing Information](#), including **Boxed WARNING**.

Important Safety Information (cont.)

Warnings and Precautions: QT Interval Prolongation

- NUPLAZID prolongs the QT interval. The use of NUPLAZID should be avoided in patients with known QT prolongation or in combination with other drugs known to prolong QT interval including Class 1A antiarrhythmics or Class 3 antiarrhythmics, certain antipsychotic medications, and certain antibiotics.
- NUPLAZID should also be avoided in patients with a history of cardiac arrhythmias, as well as other circumstances that may increase the risk of the occurrence of torsade de pointes and/or sudden death, including symptomatic bradycardia, hypokalemia or hypomagnesemia, and presence of congenital prolongation of the QT interval.

Adverse Reactions: The common adverse reactions ($\geq 2\%$ for NUPLAZID and greater than placebo) were peripheral edema (7% vs 2%), nausea (7% vs 4%), confusional state (6% vs 3%), hallucination (5% vs 3%), constipation (4% vs 3%), and gait disturbance (2% vs <1%).

Drug Interactions:

- Coadministration with strong CYP3A4 inhibitors (e.g., ketoconazole) increases NUPLAZID exposure. Reduce NUPLAZID dose to 10 mg taken orally as one tablet once daily.
- Coadministration with strong or moderate CYP3A4 inducers reduces NUPLAZID exposure. Avoid concomitant use of strong or moderate CYP3A4 inducers with NUPLAZID.

Please read the full [Prescribing Information](#), including **Boxed WARNING**, also available at NUPLAZIDhcp.com.

Reference: 1. Acadia Pharmaceuticals Inc. Nuplazid® [package insert] San Diego, CA; 2020.



©2021 Acadia Pharmaceuticals Inc. Acadia and NUPLAZID are registered trademarks and Acadia Connect is a trademark of Acadia Pharmaceuticals Inc. All rights reserved. NU-1917-v2-C 03/21.

ONCE-DAILY
NUPLAZID®
(pimavanserin) 34mg capsules